

John Hellerstedt, M.D. Commissioner

☑ CHECKLIST FOR YOUTH SPORTS FAMILIES

Page 1 of 2

A note about children and COVID-19:

The virus that causes COVID-19 can infect people of all ages. While the risk of serious illness or loss of life is greatest in those 65 years of age or older with pre-existing health conditions, persons in every age group can become infected with COVID-19 and some may become seriously ill or even die.

We should all be thankful that, with rare exceptions, COVID-19 is not claiming the lives of our children. However, we can never forget that a child with a mild or even asymptomatic case of COVID-19 can spread that infection to others who may be far more vulnerable.

COVID-19 is spread from person to person through contact that is close enough to share droplets generated by coughing, sneezing, speaking, and even just breathing. COVID-19 can also be spread by touching objects where contaminated droplets have landed. Because of this easy manner of transmission, an infant, child or young person who is infected with COVID-19 can spread the infection to others they come in close contact with, such as members of their household, teachers, or other caregivers. We have learned that infected persons with mild or even no symptoms can spread COVID-19.

These facts are vitally important when considering engaging in youth sporting activities.

One thing is for certain: We must find ways to protect our children from COVID-19 and ensure that they do not bring the infection to others, such as other household members, who may be at high risk for severe infection or even loss of life.

For adults in the workplace or other public spaces, we are confident that if recommended measures such as cloth face coverings or non-medical grade masks, respiratory etiquette, frequent hand washing / hand sanitation and environmental cleaning and sanitizing are widely observed, we can then improve safety.

However, such recommended protective measures that we can expect from adults are, for a variety of reasons, simply not possible for infants, children and youth to practice in sporting activities.

All of these factors mean that while certain precautions against the spread of COVID-19 can and will be applied to youth sports, the infection control measures that can be put in place in these settings will differ from those that are suitable for other social, business and commercial settings.

Every adult who is responsible for providing care for youth in these settings must be aware of these facts and be willing to comply with the infection control measures that will be in place in these settings. Parents should monitor the health of their children and not send them to participate in sporting activities if they exhibit any symptom of COVID-19. They should seek COVID-19 testing promptly and report results to the program given the implications for other children, families, and staff. Individuals aged 65 or older are at a higher risk of COVID-19. Parents should protect any vulnerable persons who are members of the same household or come into frequent, close contact with individuals who participate in youth sports.

About health recommendations:

Individuals age 10 or older are strongly encouraged to wear face coverings over the nose and mouth wherever it is not feasible to maintain 6 feet of social distancing from another person not in the same household.



John Hellerstedt, M.D. Commissioner

YOUTH SPORTS FAMILIES: Page 2 of 2

The following are the health recommendations for all individuals participating in youth sports. These health recommendations are not a limit on the health protocols that individuals may adopt. Individuals are encouraged to adopt additional protocols consistent with their specific needs and circumstances to help protect the health and safety of all Texans.

The virus that causes COVID-19 can be spread to others by infected persons who have few or no symptoms. Even if an infected person is only mildly ill, the people they spread it to may become seriously ill or even die, especially if that person is 65 or older with pre-existing health conditions that place them at higher risk. Because of the hidden nature of this threat, everyone should rigorously follow the practices specified in these recommendations. The virus that causes COVID-19 is still circulating in our communities. We should continue to observe practices that protect everyone, including those who are most vulnerable.

Please note, public health guidance cannot anticipate every unique situation. Individuals should stay informed and take actions based on common sense and wise judgment that will protect health and support economic revitalization.

Health recommendations for participants:				
		en the participant before attending the or worsening signs or symptoms of pos Cough Shortness of breath or difficulty breathing Chills		rting event or practice for any of the following D-19: Sore throat Loss of taste or smell Diarrhea Feeling feverish or a measured temperature
		Repeated shaking with chills		greater than or equal to 100.0 degrees Fahrenheit
		Muscle pain		Known close contact with a person who is lab
		Headache		confirmed to have COVID-19
	Carry and use hand sanitizer regularly, including before and after every meal and activity.			
Health recommendations for parents or guardians:				
	Individuals should, to the extent possible, minimize in-person contact with others not in the individual's household. Minimizing in-person contact includes maintaining 6 feet of separation from individuals. When maintaining 6 feet of separation is not feasible, other methods should be utilized to slow the spread of COVID-19, such as washing or sanitizing hand frequently, and avoiding sharing utensils or other common objects.			